

# Anterior Cruciate Ligament Injuries

## Introduction

## Physiotherapy in Kitchener for Knee



**Welcome to Kitchener Physiotherapy & Wellness patient resource about from Anterior Cruciate Ligament Injuries.**

The anterior cruciate ligament (ACL) is probably the most commonly injured ligament of the knee. In most cases, the ligament is injured by people participating in athletic activity. As sports have become an increasingly important part of day-to-day life over the past few decades, the number of ACL injuries has steadily increased. This injury has received a great deal of attention from orthopedic surgeons over the past 15 years, and very successful operations to reconstruct the torn ACL have been invented.


**This article will help you understand:**

- where in the knee the ACL is located
- how an ACL injury causes problems
- how doctors treat the condition

## **Anatomy**


**Where is the ACL, and what does it do?**

*Ligaments* are tough bands of tissue that connect the ends of bones together. The ACL is located in the center of the knee joint where it runs from the backside of the *femur* (thighbone) to connect to the front of the *tibia* (shinbone).




The ACL runs through a special notch in the femur called the *intercondylar notch* and attaches to a special area of the tibia called the tibial spine.

## Tibial Spine



The ACL is the main controller of how far forward the tibia moves under the femur. This is called *anterior translation of the tibia*. If the tibia moves too far, the ACL can rupture. The ACL is also the first ligament that becomes tight when the knee is straightened. If the knee is forced past this point, or hyperextended, the ACL can also be torn.



Other parts of the knee may be injured when the knee is twisted violently, as in a clipping injury in football. It is not uncommon to also see a tear of the *medial collateral ligament* (MCL) on the inside edge of the knee, and the *lateral meniscus*, which is the U-shaped cushion between the outer half of the tibia and femur bones.

Related Document: [Kitchener Physiotherapy & Wellness Guide to Knee Anatomy](#)

## **Knee Anatomy Introduction**

### **Causes**

#### **How do ACL injuries occur?**

The mechanism of injury for many ACL ruptures is a sudden *deceleration* (slowing down or stop), *hyperextension*, or *pivoting* in place. Sports-related injuries are the most common.

The types of sports that have been associated with ACL tears are numerous. Those sports requiring the foot to be planted and the body to change direction rapidly (such as basketball) carry a high incidence of injury. In this way, most ACL injuries are considered *noncontact*. However, contact-related injuries can result in ACL tears. For example, a blow to the outside of the knee when the foot is planted is the most likely contact-related injury.

Football is also frequently the source of an ACL tear. Football combines the activity of planting the foot and rapidly changing direction and the threat of bodily contact. Downhill skiing is another frequent source of injury, especially since the introduction of ski boots that come higher up the calf. These boots move the impact of a fall to the knee rather than the ankle or lower leg. An ACL injury usually occurs when the knee is forcefully twisted or hyperextended while the foot remains in contact with the ground. Many patients recall hearing a loud pop when the ligament is torn, and they feel the knee give way.

The number of women suffering ACL tears has dramatically increased. This is due in part to the rise in women's athletics. But studies have shown that female athletes are two to four times more likely to suffer ACL tears than male athletes in the same sports.

Recent research has shown several factors that contribute to women's higher risk of ACL tears. Women athletes seem less able to tighten their thigh muscles to the same degree as men. This means women don't get their knees to hold as steady, which may give them less knee protection during heavy physical activity. Also, tests show that women's quadriceps and hamstring muscles work differently than men's. Women's quadriceps muscles (on the front of the thigh) work extra hard during knee-bending activities. This pulls the tibia forward, placing the ACL at risk for a tear.

Meanwhile, women's hamstring muscles (on the back of the thigh) respond more slowly than in men. The hamstring muscles normally protect the tibia from sliding too far forward. Women's sluggish hamstring response may allow the tibia to slip forward, straining the ACL. Other studies suggest that women's ACLs may be weakened by the effects of the female hormone *estrogen*. Taken together, these factors may explain why female athletes have a higher risk of ACL tears.

## Symptoms

### What does a torn ACL feel like?

The symptoms following a tear of the ACL can vary. Some patients report hearing and/or feeling a *pop*. Usually, the knee joint swells within a short time following the injury. This is due to bleeding into the knee joint from torn blood vessels in the damaged ligament. The instability caused by the torn ligament leads to a feeling of insecurity and giving way of the knee, especially when trying to change direction on the knee. The knee may feel like it wants to slip backwards. There may be activity-related pain and/or swelling. Walking downhill or on ice is especially difficult. And you may have trouble coming to a quick stop.


The pain and swelling from the initial injury will usually be gone after two to four weeks, but the knee may still feel unstable. The symptom of instability and the inability to trust the knee for support are what require treatment. Also important in the decision about treatment is the growing realization by orthopedic surgeons that long-term instability leads to early arthritis of the knee.

Related Document: [Kitchener Physiotherapy & Wellness Guide to Osteoarthritis of the Knee](#)

## Diagnosis

When you visit Kitchener Physiotherapy & Wellness, our physiotherapist will first take a history and do a physical exam. The history and physical examination are probably the most important ways to diagnose a ruptured or deficient ACL.

## Bleeding into joint capsule



©MMG 2000

In the acute (sudden) injury, the swelling is a good indicator. A good rule of thumb that orthopedic surgeons use is that any tense swelling that occurs within two hours of a knee injury usually represents blood in the joint, or a hemarthrosis. If the swelling occurs the next day, the fluid is probably from the inflammatory response.

During the physical examination, special stress tests are performed on the knee. Three of the most commonly used tests are the *Lachman* test, the *pivot-shift* test, and the *anterior drawer* test. Our physiotherapist will place your knee and leg in various positions and then apply a load or force to the joint. Any excess motion or unexpected movement of the tibia relative to the femur may be a sign of ligament damage and insufficiency.

Another way to check for anterior tibial translation is with the *KT-1000* and *KT-2000 arthrometers*. The patient's leg is bent and supported on a wedge with the knee in 30 degrees of flexion. The arthrometer is placed against the knee to be tested and strapped to the lower leg. Usually, the normal knee is tested first. The arthrometer applies an anterior force of 15 pounds against the tibia. The amount of anterior tibial translation is measured. The test is repeated with a force of 20 pounds. A third test applies a manual maximal force to the posterior (back) of the tibia. This is similar to the Lachman test.

The results of these tests will help our physiotherapist determine how badly the ACL was injured. We may also combine other tests with tests of ACL integrity to determine whether other knee ligaments, joint capsule, or joint cartilage have also been injured.

Some patients may be referred to a doctor for further diagnosis. Once your diagnostic examination is complete, the physiotherapists at Kitchener Physiotherapy & Wellness have treatment options that will help

speed your recovery, so that you can more quickly return to your active lifestyle.

Kitchener Physiotherapy & Wellness provides physiotherapist services in Kitchener.

## **Our Treatment**

### **Non-surgical Rehabilitation**

When you begin your Kitchener Physiotherapy & Wellness program, our initial treatments for an ACL injury will focus on decreasing pain and swelling in the knee. We may recommend rest and mild pain medications, such as acetaminophen (Tylenol), to help decrease your symptoms.

You may need to use crutches until you can walk without a limp. Most of our ACL reconstruction patients are instructed to put a normal amount of weight down while walking. Our physiotherapist will treat swelling and pain with the use of ice, electrical stimulation, and rest periods with your leg supported in elevation.

Our physiotherapist may apply treatments such as electrical stimulation and ice to reduce pain and swelling. We then gradually add exercises to improve knee range of motion and strength to help you regain normal movement of joints and muscles.

Our physiotherapist will have you begin range-of-motion exercises right away, with the goal of helping you swiftly regain full movement in your knee. This may include the use of a stationary bike, gentle stretching, and careful pressure applied to the knee by the physiotherapist. We will also give you exercises to improve the strength of your hamstring and quadriceps muscles. As your symptoms ease and strength improves, we will guide you in specialized exercises to improve knee stability.

Our physiotherapist may suggest use of an ACL brace. This type of brace is usually custom-made and not the type you can buy at the drugstore. It is designed to improve knee stability when the ACL doesn't function properly.

We often recommend an ACL brace when the knee is unstable and surgery is not planned. As mentioned, a torn ACL that isn't corrected often leads to early knee arthritis. There is no evidence that an ACL brace will prevent further damage to the knee due to wear and tear arthritis. The ACL brace may help keep your knee from giving way during moderate activity. However, it can give a false sense of security and won't always protect the knee during sports that require heavy cutting, jumping, or pivoting. Our physiotherapists will often recommend wearing a brace for at least one year after a surgical reconstruction, so even if you decide to have ACL surgery, a brace is probably a good investment.

Although the time required for recovery varies, nonsurgical rehabilitation for a torn ACL typically lasts six to eight weeks. You can return to your sporting activities when your quadriceps and hamstring muscles are back to nearly their full strength and control, you are not having swelling that comes and goes, and you aren't having problems with the knee giving way.

### **Post-surgical Rehabilitation**

If you undergo surgery, you will probably be involved in a Kitchener Physiotherapy & Wellness progressive rehabilitation program for about four to six months after surgery to ensure the best result from your ACL reconstruction. At first, expect to see our physiotherapist about two to three times a week. If your surgery and rehabilitation go as planned during the first six weeks, you may only need to do a home program and see our physiotherapist every few weeks over the four to six month period.


At Kitchener Physiotherapy & Wellness, our goal is to help speed your recovery so that you can more quickly return to your everyday activities. When your recovery is well under way, regular visits to our office will end. Although we will continue to be a resource, you will be in charge of doing your exercises as part of an ongoing home program.

Kitchener Physiotherapy & Wellness provides services for physiotherapy in Kitchener.

## Physician Review

If there is fluid associated with your ACL injury, your doctor may need to place a needle in the swollen joint and aspirate (drain as much fluid as possible) the give relief from the swelling. The procedure also provides useful information to your doctor. If blood is found when draining the knee, there is about a 70 percent chance it represents a torn ACL. This fluid can also show if the cartilage on the surface of the knee joint was injured.

## Aspiration




Your doctor may order X-rays of the knee to rule out a fracture. Ligaments and tendons do not show up on X-rays, but bleeding into the joint can result from a fracture of the knee joint, or when portions of the joint surface are chipped off.


Magnetic Resonance Imaging is probably the most accurate test for diagnosing a torn ACL without actually looking into the knee. The MRI machine uses magnetic waves rather than X-rays to show the soft tissues of the body. This machine creates pictures that look like slices of the knee. The pictures show the anatomy, and any injuries, very clearly. This test does not require any needles or special dye and is painless.



# Magnetic Resonance Imaging (MRI)



In some cases, *arthroscopy* may be used to make the definitive diagnosis if there is a question about what is causing your knee problem.



Arthroscopy is an operation that involves inserting a small fiber-optic TV camera into the knee joint, allowing the orthopedic surgeon to look at the structures inside the joint directly. The vast majority of ACL tears are diagnosed without resorting to this type of surgery, though arthroscopy is sometimes used to repair a torn ACL.

## **Surgery**

If the symptoms of instability are not controlled by a brace and rehabilitation program, then surgery may be suggested. The main goal of surgery is to keep the tibia from moving too far forward under the femur bone and to get the knee functioning normally again.

Even when surgery is needed, most surgeons will have their patients attend physiotherapy for several visits before the surgery. This is done to reduce swelling and to make sure you can straighten your knee completely. This practice also reduces the chances of scarring inside the joint and can speed recovery after surgery.


## **Arthroscopic Method**

Most surgeons now favor reconstruction of the ACL using a piece of tendon or ligament to replace the torn ACL. This surgery is most often done with the aid of the arthroscope. Incisions are usually still required around the knee, but the surgery doesn't require the surgeon to open the joint. The arthroscope is used to view the inside of the knee joint as the surgeon performs the work. Most ACL surgeries are now done on an outpatient basis, and many patients go home the same day as the surgery. Some patients stay one or two nights in the hospital if necessary.


## **Patellar Tendon Graft**

One type of graft used to replace the torn ACL is the patellar tendon. This tendon connects the kneecap (*patella*) to the tibia. The surgeon removes a strip from the center of the ligament to use as a replacement for the torn ACL.

## **Patellar Tendon**



**Middle third of patellar ligament cut and removed**



©MMG 2000

Related Document: [Kitchener Physiotherapy & Wellness Guide to Patellar Tendon Graft Reconstruction of the ACL](#)


## Hamstring Tendon Graft

Surgeons also commonly use a hamstring graft to reconstruct a torn ACL. This graft is taken from one of the hamstring tendons that attaches to the tibia just below the knee joint. The hamstring muscles run down the back of the thigh. Their tendons cross the knee joint and connect on each side of the tibia. The graft used in ACL reconstruction is taken from the hamstring tendon, called the *semitendinosus*. This tendon runs along the inside part of the thigh and knee. Surgeons also commonly include as part of the hamstring graft a tendon just next to the *semitendinosus*, called the *gracilis*.

## Hamstring Graft




**Parts of semitendinosus and gracilis tendons removed**



©MMG 2000

When arranged into three or four strips, the hamstring graft has nearly the same strength as a patellar tendon graft.

**Tendons folded and sutured**



©MMG 2000

Related Document: [Kitchener Physiotherapy & Wellness Guide to Hamstring Tendon Graft Reconstruction of the ACL](#)

# Allograft Reconstruction

Other materials are also used to replace the torn ACL. In some cases, an *allograft* is used. An allograft is tissue that comes from someone else. This tissue is harvested from tissue and organ donors at the time of death and sent to a tissue bank. The tissue is checked for any type of infection, sterilized, and stored in a freezer. When needed, the tissue is ordered by the surgeon and used to replace the torn ACL. The allograft (your surgeon's choice of graft) can be from the tibialis tendon, patellar tendon, hamstring tendon, or *Achilles tendon* (the tendon that connects the calf muscles to the heel).

Many surgeons use patellar tendon allograft tissue because the tendon comes with the original bone still attached on each end of the graft (from the patella and from the tibia). This makes it easier to fix the allograft in place.

The advantage of using an allograft is that the surgeon does not have to disturb or remove any of the normal tissue from your knee to use as a graft. The operation also usually takes less time because the graft does not need to be harvested from your knee.

Portions of this document copyright MMG, LLC.